


National Oat Breeding Program



Contact: Pamela, Sue or Peter
Ph: 0401 122 103
0421 615 777
0401 122 127

Hay Oats - WA



- Brusher[Ⓛ]**
- Excellent hay yield in low to medium rainfall areas
 - Improved hay quality and disease resistance compared to Wintaroo
 - Flexibility in hay cutting time

- Mulgara [Ⓛ]**
- Wintaroo maturity with improved disease resistance and hay quality
 - Good grain yield and quality with high protein and low screenings
 - Care with plant density required due to large grain size

- Tungoo[Ⓛ]**
- Kangaroo replacement with improved hay quality
 - Broad spectrum of disease resistance
 - Only variety available with red leather leaf resistance

- Forester[Ⓛ]**
- Suited to high rainfall and irrigation
 - Excellent early vigour and foliar disease resistance
 - Seed available from AGF Seeds

For more information please go to www.aexco.com.au

DISEASE PROFILE

Variety	Stem rust ¹	Leaf rust ¹	BYDV ¹	Septoria ¹	Bacterial blight ¹	CCN R ¹	CCN T ²	Red leather leaf ¹	Stem diameter ³
Brusher	MR-S	R-MS	MR-MS	S-VS	MR-MS	R	MI	MS	M
Carrolup	MS	S	MS	S-VS	MR-S	S	I	S	M
Durack	MR-MS	R-S	MS-S	S-VS	MR-S	R	M/MT	MS	M
Forester	R	R-MS	MS	MS-S	MS-S	MS	MI	R-MR	MT
Kangaroo	R-S	MS-S	MR-S	MS-S	MR-MS	R	MT	MS	MF
Mulgara	MR-MS	MR	MS-S	MR-S	MR	R	MT	MS	M
Tammar	R-MR	R-MR	MS-S	MS	MR	MR	MT	R-MS	MF
Tungoo	MR-S	R-MS	MR-MS	MS-S	MR	R	MT	R	M
Walleraroo	MS-S	VS	MS	S-VS	S	R	MT	MS	F
Wandering	MR	VS	MS-S	S-VS	MR-S	VS	I	MS	M
Williams	MR	R	MR-MS	MS	R	S	I	MS	MT
Winjardie	MR-S	S-VS	MS-S	S-VS	S	S	I	MS	M
Wintaroo	MR	S-VS	MR-MS	MS-S	MR	R	MT	MS	M
Yallara	MR-MS	R	MR-MS	MS-S	MR-MS	R	I	MS	MF

¹ Disease reactions where R= resistant, MR=moderately resistant, MS=moderately susceptible, S= susceptible, VS=very susceptible

² T=tolerant, MT= moderately tolerant, MI=moderately intolerant, I=intolerant ³ F=fine, MF=moderately fine, MT=moderately thick, T=thick, VT=very thick

Note: Stem rust, leaf rust, BYDV & Septoria reactions are from WA trials, Bacterial blight, CCN & red leather leaf reactions are from SA trials

(Rust and BYDV reactions may vary in different regions and with different seasonal conditions depending on the prevalent pathotype/serotype. Monitoring your oat crop is therefore essential.)



Department of
Primary Industries and
Regional Development

